

FINITE TEMPERATURE STUDY OF NUCLEAR MATTER IN SU(2) CHIRAL SIGMA MODEL

P. K. Jena* and L. P. Singh⁺

Department of Physics, Utkal University, Vanivihar,
Bhubaneswar-751004, India.

Abstract

We study here the equation of state of symmetric nuclear matter at finite temperatures using a modified SU(2) Chiral Sigma model. The effect of temperature on effective mass, pressure, entropy and binding energy is discussed. The liquid-gas phase transition is investigated and the critical values of temperature, density and pressure are calculated.

PACS Nos: 26.60+c, 21.30.Fe, 21.60.Jz.

*email: pkjena@iopb.res.in

⁺email: lambodar@iopb.res.in

1 Introduction

Accurate description of properties of hot, dense nuclear matter is a fundamental problem in theoretical physics. To determine the properties of nuclear matter as functions of density and temperature, one must study the ground state and excited states of normal nuclei along with the highly excited states of normal nuclei created in nucleus-nucleus collisions and in nuclei far from stability, which may be created in radioactive beams[1]. As the equation of state(EOS) describes how the energy density and pressure vary with density and temperature, it is possible to study the liquid-gas phase transition, which may occur in the warm and dilute matter produced in heavy-ion collisions. Several authors had studied the liquid-gas phase transition using non-relativistic theory[2] and the critical temperature has been estimated to be in the range 15-20 MeV. But the experimental results for warm and dilute nuclear matter produced in heavy-ion collisions support a small liquid-gas phase region with a low critical temperature $T_c \approx 13.1 \pm 0.6$ MeV[3]. For symmetric nuclear matter, using relativistic mean field theory, the critical temperature given by Walecka model is $T_c \approx 18.3$ MeV which reduces to $T_c \approx 14.2$ MeV[4] if the non linear terms are included.

In the present paper, we continue our investigation with modified SU(2) chiral sigma model(MCH)[5,6] using relativistic mean-field calculations for symmetric hot, dense nuclear matter. The earlier studies have SU(2) chiral sigma model as a viable microscopic model for description of dense nuclear matter. In the present work, We study the effect of temperature on the EOS, effective mass, entropy and binding energy for symmetric nuclear matter and also investigate the liquid-gas phase transition.

This paper is organized as follows: In sec.2, we present nuclear matter at finite temperature. The results and discussion are presented in sec.3. We end with concluding remarks in sec.4.

2 Nuclear matter at finite temperature

We use MCH[5,6] model for symmetric nuclear matter since chiral model has been very successful, as such, in describing high density nuclear matter. The importance of chiral symmetry[7] in the study of nuclear matter was first emphasized by Lee and Wick[8] and has become over the years, one of most useful tools to study high density nuclear matter at the microscopic level. The nonlinear terms in the chiral sigma model give rise to the three-body forces which become significant in the high density regime[9]. Further, the energy per nucleon at saturation needed the introduction of isoscalar field[10] in addition to the scalar field of pions[11]. The MCH model[5,6] considered by us includes two extra higher order scalar field interaction terms which ensures an appropriate incompressibility of symmetric nuclear matter at saturation density. Further, the equation of state(EOS) derived from this model is compatible with that inferred from recent heavy-ion collision data[12].

The EOS for hadronic phase is calculated by using the Lagrangian density[5] (with $\hbar = c = K_{Boltzmann} = 1$),

$$L = \frac{1}{2}(\partial_\mu \vec{\pi} \cdot \partial^\mu \vec{\pi} + \partial_\mu \sigma \partial^\mu \sigma) - \frac{1}{4}F_{\mu\nu}F_{\mu\nu} - \frac{\lambda}{4}(x^2 - x_0^2)^2 - \frac{\lambda B}{6m^2}(x^2 - x_0^2)^3 - \frac{\lambda C}{8m^4}(x^2 - x_0^2)^4 - g_\sigma \bar{\psi}(\sigma + i\gamma_5 \vec{\tau} \cdot \vec{\pi})\psi + \bar{\psi}(i\gamma_\mu \partial^\mu - g_\omega \gamma_\mu \omega^\mu)\psi + \frac{1}{2}g_\omega^2 x^2 \omega_\mu \omega^\mu \quad (1)$$

In the above Lagrangian, $F_{\mu\nu} \equiv \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$ and $x = (\vec{\pi}^2 + \sigma^2)^{1/2}$, ψ is the nucleon isospin doublet, $\vec{\pi}$ is the pseudoscalar-isovector pion field, σ is the scalar field and ω_μ , is a dynamically generated isoscalar vector field, which couples to the conserved baryonic current $j_\mu = \bar{\psi}\gamma_\mu\psi$. B and C are constant coefficients associated with the higher order self-interactions of the scalar field.

The masses of the nucleon, the scalar meson and the vector meson are respectively given by

$$m = g_\sigma x_0, \quad m_\sigma = \sqrt{2\lambda}x_0, \quad m_\omega = g_\omega x_0 \quad (2)$$

Here x_0 is the vacuum expectation value of the σ field, g_ω and g_σ are the coupling constants for the vector and scalar fields respectively and $\lambda = (m_\sigma^2 - m_\pi^2)/(2f_\pi^2)$, where

m_π is the pion mass , f_π is the pion decay coupling constant .

Using Mean-field approximation, the equation of motion for vector meson field is

$$\omega_0 = \frac{n_B}{g_\omega x^2} \quad (3)$$

n_B is the baryon number density at temperature T and is given by

$$n_B = \frac{\gamma}{(2\pi)^3} \int_0^\infty d^3k [n(T) - \bar{n}(T)] \quad (4)$$

with $n(T) = \frac{1}{e^{(E^*(k)-\nu)\beta+1}}$, $\bar{n}(T) = \frac{1}{e^{(E^*(k)+\nu)\beta+1}}$,

where $E^*(k) = (k^2 + y^2 m^2)^{1/2}$, the effective nucleon energy, $\nu = \mu_B - C_\omega n_B / y^2$, the effective baryon chemical potential, $\beta = 1/K_B T$, γ is the spin degeneracy factor with $n(T)$ and $\bar{n}(T)$ being Fermi-Dirac distribution functions for particle and anti-particle at finite temperature and $y \equiv x/x_0$ is the effective mass factor which must be determined self consistently from the equation of motion for scalar field which is given by

$$(1-y^2) - \frac{B}{m^2 C_\omega} (1-y^2)^2 + \frac{C}{m^4 C_\omega^2} (1-y^2)^3 + \frac{2C_\sigma C_\omega n_B^2}{m^2 y^4} - \frac{C_\sigma \gamma}{\pi^2} \int_0^\infty \frac{dk k^2 (n + \bar{n})}{\sqrt{k^2 + m^{*2}}} = 0 \quad (5)$$

$m^* \equiv ym$ is the effective mass of the nucleon and the coupling constants are

$$C_\sigma \equiv \frac{g_\sigma^2}{m_\sigma^2}, \quad C_\omega \equiv \frac{g_\omega^2}{m_\omega^2} \quad (6)$$

The energy density and pressure at finite temperature and finite density are given by

$$\begin{aligned} \epsilon &= \frac{m^2(1-y^2)^2}{8C_\sigma} - \frac{B}{12C_\omega C_\sigma} (1-y^2)^3 + \frac{C}{16m^2 C_\omega^2 C_\sigma} (1-y^2)^4 \\ &\quad + \frac{C_\omega n_B^2}{2y^2} + \frac{\gamma}{2\pi^2} \int_0^\infty dk k^2 \sqrt{(k^2 + m^{*2})} (n + \bar{n}) \\ P &= -\frac{m^2(1-y^2)^2}{8C_\sigma} + \frac{B}{12C_\omega C_\sigma} (1-y^2)^3 - \frac{C}{16m^2 C_\omega^2 C_\sigma} (1-y^2)^4 \\ &\quad + \frac{C_\omega n_B^2}{2y^2} + \frac{\gamma}{6\pi^2} \int_0^\infty \frac{dk k^4 (n + \bar{n})}{\sqrt{(k^2 + m^{*2})}} \end{aligned} \quad (7)$$

The entropy density can be obtained as

$$S = \frac{1}{T} \left[\frac{C_\omega n_B^2}{y^2} + \frac{\gamma}{2\pi^2} \int_0^\infty dk k^2 \sqrt{(k^2 + m^{*2})} (n + \bar{n}) + \frac{\gamma}{6\pi^2} \int_0^\infty \frac{dk k^4 (n + \bar{n})}{\sqrt{(k^2 + m^{*2})}} \right] - \frac{\mu_B n_B}{T} \quad (8)$$

The values of four parameters C_σ, C_ω, B and C occurring in the above equations are obtained by fitting with the saturation values of binding energy/nucleon (-16.3 MeV), the saturation density (0.153 fm^{-3}), the effective(Landau) mass ($0.85M$)[13], and nuclear incompressibility ($\sim 300 \text{ MeV}$), in accordance with recent heavy-ion collision data[12] are $C_\omega = 1.999 \text{ fm}^2$, $C_\sigma = 6.816 \text{ fm}^2$, $B = -99.985$ and $C = -132.246$.

Thus the energy density, pressure and entropy density for symmetric nuclear matter at finite temperature and finite density can be evaluated by first solving equations (4) and (5) self consistently for fixed choosen value of the inverse temperature β with variation of n_B to get μ_B and y and then making substitution in equations (7) and (8).

3 Results and discussions

In Fig.1, we show pressure as a function of density n_B at different temperatures. For a given n_B , the pressure has the usual trend of increasing with temperature[4]. As the temperature increases the EOS becomes stiffer. Pressure has a non-zero value for $n_B = 0$ at and above a temperature of 200 MeV. It indicates that pressure has a contribution arising from the thermal distribution functions for baryons and anti-baryons as well as from the non-zero value of the scalar field. Similar results were also obtained by Panda *et al.*[14] for symmetric nuclear matter. The non-zero value for scalar σ field has also been observed in Walecka model[4].

Fig.2 shows the behavior of the effective nucleon mass with density at different temperatures. We observe that for low temperatures the results are not so different from those obtained at zero temperature. This implies that for low temperatures, the density dependence is more important than the temperature dependence. At 200 MeV the effective nucleon mass m^* first increases and then decreases slowly. This behavior indicates that the effective nucleon mass is considerable in contrast to the Walecka model[15] but quite similar to that in ZM models[16].

In Fig.3 we show the effective mass as a function temperature at zero density($n_B = 0$) and saturation density($n_B = 0.153 \text{ fm}^{-3}$). It is observed that for $n_B = n_0$, the M^*




Figure 1: Pressure (P) as function of baryon number density at different temperatures.

first increases slowly and then falls suddenly at about $T \approx 240$ MeV. But for $n_B = 0$, M^* remains almost constant as the temperature increases and falls suddenly at about $T \approx 235$ MeV. This result shows that a first order phase transition appears at $n_B = 0$, $T \approx 235$ MeV which is similar to the result obtained for Walecka model, which has such a phase transition at $T \approx 185$ MeV[17]. Because of strong strong attraction between the nucleons at high temperatures the nucleon antinucleon pairs can be formed which may lead to abrupt change in M^* to take place in high temperature region.

The entropy density as a function of density at different temperatures is presented in Fig.4. It is observed that entropy density is non-zero even at vanishing baryon density at and above a temperature of 150 MeV with contributions from the non-zero value of the sigma field. Similar behavior was also observed for entropy density in the Walecka




Figure 2: *Effective mass as function of baryon number density at different temperatures.*

model and ZM model calculations[16]. This increase of entropy density with increase of temperature indicates a phase transition.

Fig.5 shows the energy per nucleon as a function of the baryon density at different temperatures. With the increase of temperature the minimum shifts towards higher densities and for higher temperatures the minimum of the curve becomes positive. It is also observed that as the temperature increases, the nuclear matter becomes less bound and the saturation curves in the MCH model are flatter than those observed in Walecka model[15,16]. This result implies that the nuclear matter EOS in MCH model is softer than that obtained in Walecka model.

Lastly we present the pressure of nuclear matter as a function of baryon density for low temperatures in Fig.6. The figure shows that at zero temperature, the pressure first




Figure 3: *Effective mass as function of temperature for constant baryon number densities.*

decreases, then increases and passes through $P = 0$ at $n_B = n_0$ (saturation density), where the binding energy per nucleon is a minimum. Decrease of pressure with density implies a negative incompressibility, $K = 9(\frac{\partial P}{\partial n_B})$, which is a sign of mechanical instability. When the temperature increases the region of mechanical instability decreases and disappears at the critical temperature T_c , which is determined by $\frac{\partial P}{\partial n_B} \big|_{T_c} = \frac{\partial^2 P}{\partial^2 n_B} \big|_{T_c} = 0$, above which the liquid-gas phase transition is continuous. We have obtained the value of critical temperature $T_c \approx 17.2 \text{ MeV}$, critical density $n_c \approx 0.045 \text{ fm}^{-3}$, critical pressure $p_c \approx 0.274 \text{ MeV/fm}^{-3}$ and which is in fair agreement with the results obtained in other studies [15, 16, 18]




Figure 4: *Entropy density as function of baryon number density at different temperatures.*

4 Conclusions

In this work, we have extended our earlier[6] investigations with MCH model at zero temperature to study the thermodynamic properties of symmetric nuclear matter at finite temperature. We have presented variations of pressure, effective nucleon mass, entropy density and energy per nucleon with respect to density for various temperatures. At zero density it is found that this model exhibit a phase transition at $T \approx 235$ MeV just as obtained in the Walecka model at $T \approx 185$ MeV[17]. We find in the zero temperature case, the MCH model gives a softer EOS of nuclear matter at finite temperature than the Walecka model. We also find that the model under investigation indicates a liquid-gas phase transition and the critical temperature is found to be $T_c \approx 17.2$ MeV above




Figure 5: *Energy per nucleon as function of baryon number density at different temperatures.*

which the liquid gas phase transition is continuous. The fair amount of agreement of our results with those of similar studies undertaken with different models strongly support MCH model as a viable model for description of nuclear matter at high density.

Acknowledgements

P.K.Jena would like to thank Council of Scientific and Industrial Research, Government of India, for the award of SRF, F.No. 9/173 (101)/2000/EMR-I. We are also thankful to Institute of Physics, Bhubaneswar, India, for providing the library and computational facility.




Figure 6: Pressure (P) as function of baryon number density at different temperatures.

References

- [1] H. Müller and B. D. Serot, *Phy. Rev. C* **52**, 2072(1995).
- [2] W. A. Kupper, G. Wegmann and E. R. Hilf, *Ann. phys.(N.Y.)* **88**, 454(1974); B. Freedman and V. R. Pandharipande, *Nucl. phys. A* **361**, 502(1981); H. Jaqaman, A. Z. Mekjian and L. Zamick, *Phy. Rev. C* **27**, 2782(1983); D. Bandyopadhyay, C. Samanta, S. K. Samaddar and J. N. De, *Nucl. phys. A* **511**, 1(1990).
- [3] T. Li *et.al*, *Phy. Rev. C* **49**, 1630(1994).
- [4] R. J. Furnstahl and B. D. Serot, *Phys. Rev.C* **41**, 262(1990).
- [5] P. K. Sahu and A. Ohnishi, *Prog. Theor. phys.* **104**, 1163 (2000).

- [6] P. K. Jena and L. P. Singh, Mod. Phys. Lett. A **17**, 2633 (2002); P. K. Jena and L. P. Singh, Mod. Phys. Lett. A **18**, 2135 (2003).
- [7] M. Gell-Mann and M. Levy, Nuovo Cim. **16**, 705 (1960).
- [8] T. D. Lee and G. C. Wick, Phys. Rev. D **9**, 2241 (1974).
- [9] A. D. Jackson, M. Rho, Nucl. Phys. A **407**, 495 (1985).
- [10] J. Boguta, Phys. Lett. B **120**, 34 (1983); **128**, 19 (1983).
- [11] P. K. Sahu, R. Basu and B. Datta, Astrophys.J. **416**, 267 (1993).
- [12] P. K. Sahu, A. Hombach, W. Cassing, M. Effenberger and U. Mosel, Nucl.Phys. A **640**, 493 (1998) ; P. K. Sahu, W. Cassing, U. Mosel and A. Ohnishi, Nucl. Phys. A **672**, 376 (2000) ; P. K. Sahu, Phys. Rev. C **62**, 045801 (2000).
- [13] P.Möller, W.D.Myers, W.J.Swiatecki and J.Treiner, Atomic Data Tables **39**, 225 (1988).
- [14] P. K. Panda, A. Mishra, J. M. Eisenberg and W. Greiner Phys. Rev. C **56**, 3134(1997).
- [15] Guo Hua, Liu Bo, M. Di Toro, Phys. Rev. C **62**, 035203(2000).
- [16] M. Malheiro, A. Delfino and C. T. Coelho, Phys. Rev. C **58**, 426(1998).
- [17] J. Theis, H. Stöcker and J. Polonyi, Phys.Rev.D **28**, 2286 (1983).
- [18] P. K. Panda, G. Krein, D. P. Menezes and C. Providencia, Phys. Rev. C **68**, 015201(2003).